



PARENTS!



Protect your child from these serious diseases that can cause your child to become very ill or hospitalized. These diseases may cause blindness, deafness, brain damage, or in rare cases, death...

Hepatitis B
Diphtheria
Pertussis (Whooping Cough)
Measles
Mumps
Rubella
Polio

Tetanus
Pneumococcus
Rotavirus
Haemophilus influenzae type b
Varicella (Chickenpox)
Hepatitis A
Meningitis
Influenza (Flu)

Vaccines...

- *Protect your child against these diseases*
 - *Help your child develop immunity without getting sick*
 - *Are safe and effective*
 - *Are required for your child to attend day care and school*

“Start your baby’s shots on time and stay on schedule”

Visit your doctor, or for information on where your child can be vaccinated,

Dial 311

RECOMMENDED PRESCHOOL IMMUNIZATION SCHEDULE, JANUARY 2007

	BIRTH	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	4 YEARS
Hepatitis B HepB	√	√		√ 6-18 months			
Diphtheria, Tetanus, Pertussis DTaP		√	√	√		√ 12-15 months	√
Haemophilus influenzae type b Hib		√	√	√		√ 12-15 months	
Polio IPV		√	√			√ 6-18 months	√
Pneumococcal Disease PCV		√	√	√	√ 12-15 months		
Rotavirus RV		√ 6-12 weeks	√	√ ≤ 32 weeks			
Measles, Mumps, Rubella MMR					√		√
Varicella Var					√ 12-15 months		√
Hepatitis A HepA					2 doses 12-23 months	The 2 nd dose should be given at least 6 months after the 1 st dose.	
Influenza				Yearly for all children 6-59 months.			
Meningococcal MPSV						Children 2 years and older in certain high-risk groups.	



THE CITY OF NEW YORK
DEPARTMENT OF HEALTH & MENTAL HYGIENE
Michael R. Bloomberg Mayor Thomas R. Frieden, M.D., M.P.H. Commissioner





¡PADRES!



Proteja su niño contra estas serias enfermedades que pueden causar que su niño se enferme gravemente o sea hospitalizado. Estas enfermedades pueden causar ceguera, sordera, daño cerebral y hasta la muerte...

- | | |
|----------------------------|-------------------------------|
| Hepatitis B | Tétano |
| Difteria | Neumocócica |
| Pertussis (Tos ferina) | Rotavirus |
| Sarampión | Haemophilus influenzae tipo b |
| Papera | Varicela |
| Rubéola (Sarampión Alemán) | Hepatitis A |
| Polio | Meningitis |
| | Influenza (Gripe) |

Las vacunas...

- Protejan contra estas enfermedades
 - Ayuda a su niño desarrollar inmunidad sin la enfermedad
 - Se requieren para que su niño entre la escuela y a centros de cuidado diurno

“Comience las vacunas de su niño a tiempo y manténgalo en el itinerario.”

Visite su médico, ó para obtener información en donde vacunar a su niño,

Llame al 311

ITINERARIO PRE-ESCOLAR RECOMENDADO DE INMUNIZACIÓN, ENERO 2007

	NACIMIENTO	2 MESES	4 MESES	6 MESES	12 MESES	15 MESES	4 AÑOS
Hepatitis B HepB	√	√		√ 6-18 meses			
Difteria, Tétano, Pertussis DTaP		√	√	√		√ 12-15 meses	√
Haemophilus influenzae tipo b Hib		√	√	√		√ 12-15 meses	
Polio IPV		√	√			√ 6-18 meses	√
Neumocócica PCV		√	√	√	√ 12-15 meses		
Rotavirus RV		√ 6-12 semanas	√	√ ≤32 semanas			
Sarampión, Papera, Rubéola MMR					√		√
Varicela Var					√ 12-15 meses		√
Hepatitis A HepA					2 dosis 12-23 meses	La segunda dosis debe ser administrada seis meses después de la primera dosis	
Influenza				Anualmente para todo niños 6 a 59 meses de edad.			
Meningococcal MPSV						Niños mayores de 2 años y grupos de alto riesgo	



THE CITY OF NEW YORK
 DEPARTMENT OF HEALTH & MENTAL HYGIENE
 Michael R. Bloomberg Thomas R. Frieden, M.D., M.P.H.
 Alcalde Comisionado
nyc.gov/health

